
Utilizing High Resolution Topography in WRF-ARW and LAPS/STMAS with TopoGrabber



Brad Beechler
Presented to the LAPS Workshop Tuesday, October 26th, 2010

Motivation

- I. Orography has a major and direct impact on the surface wind's advection.
- 2. Orography also affects microphysical processes such as the release of latent heat and the formation of precipitation.
- 3. For fire weather, regional severe storm, and accurate wind farming forecasts high resolution topography information is crucial.
- 4. Future modeling efforts will expect access to this improved topography and land use data.



Description of TopoGrabber

Includes: Topo_Grabber.py: A python GUI workflow wrapper.

Topo_Grab.py :A command line program that downloads

domain specific geotiff data from multiple

online sources.

GeoTiff2WPS.x : Converts geotiff data into raw WPS files.

GeoTiff2CDF.x : Converts geotiff data into netCDF format.*

Requires: Along with a C and Fortran compiler this package needs

NetCDF library for C

Python version 2.6 or higher

lib_tiff library (included)

lib_geotiff library (included)

Environment variables pointing to the lib_geotiff librarty example: export LD_LIBRARY_PATH="/home/you/src/lib_geotiff/lib" export LD_RUN_PATH="/home/you/src/lib_geotiff/lib"

Application to WRF


Useage:

- I: Create a directory under your geogrid data root called topo_Is (one second topography)
- 2: Replace or link your GEOGRID.TBL (located under the /WPS/geogrid directory) with our GEOGRID.TBL.HRTOPO file. This will tell georgid that there is now a directory under its georgid data root called /topo_Is with your data in it.
- 3: Run the python program Topo_Grabber.py
- 4: Take the WPS raw files (eg 05001-05100.05201-05300) that were created and put them in your topo_Is directory under the geogrid data path (eg / bin/WPS/GEOG/topo_Is).
- 5: Run georgid.exe like normal.


Topo Grabber:

000		12 11 11 11 11 11			
000	X To	ppoGrabber		000	X TopoGrabber: Source Editor
Topo Data Source:	ned3 —		Add/Edit Source	Name:	ned3
Domain:	Boulder		Add/Edit Domain	Server URL:	http://extract.cr.usgs.gov/Website/distreq/RequestSummar
Latitude Range (North):	40.0895	to	39.9395	Product code:	ND302XT
Longitude Range (East):	-105.1525	to	-105.3025	Destination variable:	ZSF
GeoTIFF Output Directory:	boulderTIFF		-100.0020	Units:	meters
Geogrid Output Directory:	boulderGrid		Export single tile	Number of categories:	None
Geogra Output Directory:			-	Tile size:	1300
	Convert!		Clear Output	Description:	National Elevation Dataset 1/3 arcsecond resolution.
Output:				Delete source?	」 Yes
Wesley Smith (+\ Conversion run #1:	0(_)0000 TopoGrabber ver	Brad Beechler		Name: Center Latitude:	opoGrabber: Domain Editor Boulder 40.014500
Using data source: ned3 Domain bounds: LatMax: 40.0895 LatMin: 39.9395 LonMax: -105.152 LonMin: -105.302	5 E 5 E			Center Longitude: DX: DY: Description:	0.150000 0.150000 Boulder, CO. The nicest place on
Domain bounds: LatMax: 40.0895 LatMin: 39.9395 LonMax: -105.152	5 E 5 E			DX: DY: Description:	0.150000 0.150000 Boulder, CO. The nicest place on Earth.
Domain bounds: LatMax: 40.0895 LatMin: 39.9395 LonMax: -105.152 LonMin: -105.302	5 E 5 E ry: boulderTIFF	ri d		DX: DY:	0.150000 Boulder, CO. The nicest place on Earth. Yes
Domain bounds: LatMax: 40.0895 LatMin: 39.9395 LonMax: -105.152 LonMin: -105.302 Topo data output directo Geogrid data output dire	5 E 5 E ry: boulderTIFF ctory: boulderGr			DX: DY: Description:	0.150000 0.150000 Boulder, CO. The nicest place on Earth.


Boulder Topography 1550m – 2035m




Flatirons Topography




Boulder Land Use (different legends)




Mt. St. Helens Topography 550m - 2550m



Grand Canyon Topography 800m – 2180m





Future Work


- 1. This 1/3 arc second (roughly 30m) data is currently being upgraded to 1/9 arc second resolution.
- 2. This data is only available for the continental United States, Hawaii, Alaska, and Puerto Rico. Hopefully this will be expended soon.
- 3. There are still major bugs in the land use ingestion function having to do with using different map geometries.
- 4. There are minor bugs in the code that need to be addressed dealing with usability and portability.
- 5. GeoTiff2CDF.x needs to be written (Ingest for LAPS/STMAS)
- 6. We need to beta test this process, who wants to help?

Contact Info

To get your own copy of TopoGrabber visit:

http://laps.noaa.gov/topograbber/

You also can find the site by typing 'topograbber' into Google.



Wesley S. Smith – wssmith@willamette.edu
Brad E. Beechler – bbeechler@precisionwind.com